

Stomax-40

Stomax-40

Omeprazole for Injection BP

40 mg.



**I.V. USE ONLY
(As Lyophilized Powder)**



Composition :
Each vial contains :
Sterile Omeprazole Sodium BP
eq. to Omeprazole 40mg.

Dosage :
As directed by the Physician.

**Storage below 25°C till
reconstitution. Protect from
direct sunlight & moisture.**

Keep all medicine out of reach
of children.

Direction for use : Reconstitute with
10ml. sterile water for injections
supplied with this pack. Pick the 10ml
ampoule of sterile water for injection
only once & discard the ampoule.
The reconstituted solution should
be used immediately and not to be
stored.

Caution : If any foreign particle is
visible in the vial after dissolving
the content. Please do not use
and return for replacement.

LOSARMIN
LOSARTAN POTÁSICO
TABLETAS RECUBIERTAS 50 mg

Composición:

Cada tableta recubierta contiene:

Losartán Potásico USP 50 mg

Excipientes c.s.

- Lactosa Monohidrato

- Almidón de Maíz Pregelatinizado.

Acción Terapéutica: Antihipertensivo.

Vía de Administración: Oral.

Dosificación:

Según lo indicado por el médico.

Almacenamiento:

Almacenar por debajo de 30°C

protegido de la luz y la humedad.

Mantener el medicamento

fuera del alcance de los niños.

Por favor, lea el prospecto adjunto

cuidadosamente antes de usar.

VENTA BAJO RECETA MÉDICA.


VIA ORAL

LOSARMIN
LOSARTAN POTÁSICO

TABLETAS
RECUBIERTAS 50 mg

100
TABLETAS

LOSARMIN
LOSARTAN POTÁSICO





Rx

Chlorzoxazone & Paracetamol Tablets

Myopar
TABLETS

10 x 10 Tablets

COMPOSITION:

Each film coated tablet contains:

Chlorzoxazone	USP	250 mg
Paracetamol	IP	325 mg
Excipients		q.s.

Colour: Titanium Dioxide IP

Dosage: As directed by the Physician.

Storage: Store in cool, dark & dry place.

Protect from direct sunlight.

Keep the medicine out of reach of children.

Warning: Taking more than daily dose of Paracetamol may cause serious liver damage and allergic reactions (eg. Swelling of face, mouth & throat difficulty in breathing, itching or rash).

SCHEDULE H - PRESCRIPTION DRUG - CAUTION
Not to be used by etc. without the prescription of a
Registered Medical Practitioner.

AMPIGEN S

AMPIGEN S

**Ampicillin and Sulbactam
For Injection USP**

Powder for preparing a
solution for Injection

**FOR IM/IV USE
ONLY**

1500 mg



АМПИГЕН С

Состав:

Каждый флакон содержит:	
Ампициллин натрия (стерильный)	USP 1000mg
сульбактам натрия (стерильный)	USP 500mg

Дозировка:
По указанию врача

Хранение:
Хранить при температуре ниже 30 градусов, в
защитном от света месте. Не замораживать.


Приготовление растворов должно быть использовано сразу
после приготовления.


Внимание: Приготовление растворов не должно содержать
видимых не вооруженным глазом частиц.

Отпускается по рецепту врача

Хранить все медикаменты в
недоступном для детей месте







Napxen

Naproxen
Tablets BP 500 mg

Napxen

For Human Use.
Mfg. Lic. No.: 06 Dross 14/04/2007
Fab. Lic. N°: 06 Dross 14/04/2007
Batch No.: JDT-3216
N° de lot: 09/2025
Mfg. Date: 08/2028
Date de fabrication:
Exp. Date:
Date de expiration:

INOSMIN
Tablets

10x10 Tablets

INOSMIN

Tablets

Myo-Inositol & Metformin
Hydrochloride (SR) Tablets


Composition :
Each film coated bilayered tablet contains:
Myo-Inositol BP 600 mg
Metformin Hydrochloride BP 500 mg
(As sustained release) q.s.


Excipients

Dosage :
As directed by the Physician.


Storage :
Store in a cool, dry & dark place.
Protect from light & moisture, at a
temperature not exceeding 25°C.
Keep out of reach of children.











CEFTRIZOL
Injection

CEFTRIZOL

Injection

Ceftriaxone for Injection USP

For I.M/I.V Use Only

1000mg.

Composition:
Each vial contains:
Sterile Ceftriaxone Sodium USP
eq. to Ceftriaxone 1000mg.

Dosage:
As directed by the Physician.

Storage:
Store in a cool, dry & dark place.
Protect from light & moisture. At a
temperature not exceeding 25°C.

Keep all medicine out of reach
of children.

Direction for use: Reconstitute with
10ml. sterile water for injections
supplied with this pack. Pick the 10ml
ampoule of sterile water for injection
only once & discard the ampoule.
The reconstituted solution should
be used immediately and not to be
stored.

Caution: If any foreign particles
visible in the vial after dissolving
the content. Please do not use
and return for replacement.



COMPOSITION:
Each hard gelatin capsule
contains:
Omeprazole BP 20 mg
(As enteric coated
granules)
Excipients Q.S.
Colour: Approved colour
used in empty hard gelatin
capsule shells

DOSAGE: As directed by the
physician.
Capsules should be swallowed
whole, not to be chewed or
crushed.
STORAGE: Store below 30°C.
Protect from light & moisture.
**KEEP OUT OF REACH OF
CHILDREN**

R_x
Omeprazole Capsules 20 mg
PPSOME 20



10x10 Capsules

Omeprazole Capsules 20 mg
PPSOME 20







Composition	IP	1.0% w/w
Chlorhexidine Dihydrochloride	IP	1.0% w/w
Equivalents to Chlorhexidine Sodium	IP	1.0% w/w
Essential Oil	IP	1.0% w/w
Methyl Salicylate	IP	1.0% w/w
Perfumed	IP	1.0% w/w
Cycloheximide Hydrochloride	IP	1.0% w/w
Oil Peppermint	IP	1.0% w/w
Essential Alcohol	IP	1.0% w/w
Oil Borneo	IP	1.0% w/w

Storage 1: As directed by the Manufacturer.
Storage 2: Store in a cool, dry, dark place.
Protect from light & moisture, at a temperature not exceeding 20°C.
*Apply 1-2 times a day. Gently rub on the affected area
the first 3 days.
Puncture holes with a sharp point of card.
Replace caps after use.*

CAUTION: Not to be sold by retail without the prescription of a Registered Medical Practitioner.

FOR EXTERNAL USE ONLY

Diclofenac Diethylamine, Linseed Oil, Methyl Salicylate,
Menthol and Cyclobenzaprine Hydrochloride Gel

30 gm



The effective pain reliever



CAPSIBENZ™ Gel

Траксовер

5 ампул
по 5 мл

траксовер

500 мг/5мл

Транексамовая кислота ВР для в/в инъекции

500
МГ



При обнаружении
повреждения ампулы,
не использовать данное
препарирование.

Отпускается по рецепту

Хранение: в температуре
ниже 25°С в сухом и
защитном от света месте.
Не замораживать.
Хранить в оригинальной
упаковке для детей месте

Каждая мл содержит:
Транексамовая кислота ВР 400 мг
Стерилизованная вода для инъекций 0,5.

Дозировка:
По указанию врача





Getzee

100 ml

Cetirizine Oral Solution BP 5mg/5ml

Getzee



Composition :
Each 5 ml. contains :
Cetirizine Hydrochloride USP 5 mg
Flavoured base q.s

Dosage :
As directed by the Physician.

Storage : Store in a cool,
dry place, Protect from
light at a temperature not
exceeding 25°C.

**SHAKE WELL BEFORE
USE**

Keep the medicine out
of reach of children.
For Human use.

Manufactured by : Fabrique par :
JPEE DRUGS
Plot No. 53, Sector-84, SOUL,
Haridwar (Uttaranchal)
(WHO-GMP, G.P.A. & ISO CERTIFIED COMPANY)

Cromsee
Sodium Cromoglicate
Eye Drops BP 2% w/v

10 ml.

Cromsee

**Sodium Cromoglicate
Eye Drops BP 2% w/v**



Composition:
Sodium Cromoglicate BP 2% w/v
Benzalkonium Chloride BP 0.01 w/v
Solution
(As preservative)
Sterile isotonic base q.s

Dosage:
As directed by Physician.
**Store away from direct
heat & light & not
exceeding 25°C.**
Use the solution within
one month after opening
the container.

Warning:
If irritation Persists or
Increases discontinue the
use and consult Physician.
Do not touch the dropper tip
or other dispensing tip to any
surface since this may
contaminate solution.

GENPEX

10 ml.

GENPEX

Gentamicin

Eye/Ear Drops BP

0.3% w/v



Composition :
Gentamicin Sulfate BP
Eq. to Gentamicin 0.3% w/v
Benzalkonium Chloride
Solution BP 0.03% v/v
(As Preservative)
Sterile Isotonic
Aqueous base q.s

Dosage :
As directed by Physician.
Store away from direct
heat & light & not
exceeding 25°C.
Use the solution within
one month after opening
the container.

Warning :
If irritation Persists or
Increases discontinue the
use and consult Physician.
Do not touch the dropper tip
of other dispensing tip to any
surface since this may
contaminate solution.

Gen-D
Gentamicin &
Dexamethasone
Eye/Ear Drops

Gentamicin & Dexamethasone Eye/Ear Drops

Gen-D



10 ml

Composition:
Gentamicin Sulfate BP 0.3% w/v
Eq. to Gentamicin USP 0.01% w/v
Dexamethasone (As Dexamethasone Sodium Phosphate USP) 0.01% w/v
Benzalkonium Chloride Solution BP 0.02% w/v
(As Preservative) q.s.
Sterile Isotonic base q.s.

Dosage:
As directed by Physician.
Store away from direct heat & light & not exceeding 25°C.
Use the solution within one month after opening the container.


Warning:
If irritation Persists or Increases discontinue the use and consult Physician. Do not touch the dropper tip of other dispensing tip to any surface since this may contaminate solution.

Timojet
Timolol Eye Drop BP
0.5% w/v

10 ml.

Timojet

Timolol Eye Drop BP
0.5% w/v



Composition :
Timolol Maleate BP 0.5% w/v
Eq. to Timolol
Benzalkonium Chloride BP 0.02% v/v
Solution (As Preservative)
Sterile Isotonic base q.s

Dosage :
As directed by Physician.
Store away from direct
heat & light & not
exceeding 25°C.
Use the solution within
one month after opening
the container.

Warning :
If irritation Persists or
Increases discontinue the
use and consult Physician.
Do not touch the dropper tip
of other dispensing tip to any
surface since this may
contaminate solution.



ARIPENEM
1 gm.

1 gm.

Injection

R_x

Meropenem
Injection IP 1.0gm.

ARIPENEM

INJECTION
1 gm.

FOR I.V. USE ONLY



Composition:
Each vial contains:
Meropenem (Series) IP
Eq. to Anhydrous Meropenem 1000 mg.
Sodium Carbonate IP
anhydrous (Series) IP 90.2 mg.
Eq. to Sodium
This is not a multi dose vial.
Any unused portion must be discarded.
Any unused portion must be discarded.

Dosage: As directed by the
Physician.

Storage: Store in a cool, dry &
dark place. Protect from light
and moisture.

Keep the medicine out of reach
of children.

Dissolve the contents of the vial
in Sterile Water for injection IP
for IV administration. Use freshly
prepared solution. After resuspension
occurs on addition of Sterile water
for injection IP.

SCHEDULE H PRESCRIPTION DRUGS - CAUTION
As a Schedule H drug, this preparation is a
prescription drug and its use should be
under the strict supervision of a
Registered Medical Practitioner.

Manufactured in India by
IPSC DRUGS
Plot No-63, Sector-14, Gurgaon, Haryana (India)
(IPSC DRUGS is a subsidiary of IPSC)

ELFORD-MR
एलफोर्ड एमआर
Tablets

R_x

20x10 Tablets

Aceclofenac, Paracetamol & Chlorzoxazone Tablets

ELFORD-MR

एलफोर्ड एमआर

Tablets

ELFORD
HEALTHCARE

Composition :
Each uncoated tablet contains :
Aceclofenac P 100mg
Paracetamol P 325mg
Chlorzoxazone USP 250mg
q.s.
Excipients
Colour : Turquoise P-5F
Dosage : As directed by the Physician.
Storage : Store in a cool, dry & dark place.
Protected from light.
Keep the medicine out of reach of children.


SCHEDULE H PRESCRIPTION DRUG-CAUTION
Not to be sold by retail without the prescription
of a Registered Medical Practitioner.

Caution : Taking more than daily dose of Paracetamol
may cause serious liver damage or allergic reactions
(e.g. swelling of the face, mouth & throat, difficulty in
breathing, itching or rash).

Manufactured in India by
JPEE DRUGS
Plot No-52, Sector-64, SOCTA, Gurgaon (Haryana)
www.jpee.com and a GMP CERTIFIED COMPANY













NOURISH VITA



200ml

NOURISH VITA

Cyproheptadine HCL
with Multivitamin
Syrup



Composition:

Each 15ml contains:
Cyproheptadine Hydrochloride 8 mg equivalent to
Cyproheptadine HCl Anhydrous 5 mg
Thiamine Hydrochloride (Vitamin B1) 3 mg
Riboflavin (Vitamin B2) as Riboflavin 2.2 mg
Phosphate sodium 50 mg
Pyridoxine Hydrochloride (Vitamin B6) 5 mg
Cyanocobalamin (Vitamin B12) 5 mcg
D-Pantothene USP 15 mg
Follicles Syntex base
Dosage: As directed by the physician.
Storage: Store below 30°C, protect from light.
Keep out of reach of children.
Please read the package insert carefully
before use.
Keep the bottle cap tightly closed.

SHAKE WELL BEFORE USE